

【格式範例】

雄鼠高泌乳素血症誘發之睪酮低常與萊氏細胞的自噬作用有關

黃志賢^{1,2,3} 蔡宜庭³ 王中麟³ 江怡德^{1,3} 浦筱峰³ 林登龍^{1,2} 陳光國^{1,2}

¹台北榮民總醫院外科部泌尿外科 ²國立陽明大學醫學院泌尿學科 ³生理學科

Hyperprolactinemia-induced hypogonadism in male rats is related with autophagy of Leydig cells

William J Huang^{1,2,3}, Yi-Ting Tsai³, Zhong-Lin Wang³, Yi-Te Chiang^{1,3}, Hsiao-Fung Pu³, Alex TL Lin^{1,2}, Kuang-Kuo Chen^{1,2}

¹Division of Urology, Department of Surgery, Taipei Veterans General Hospital, Taipei, Taiwan.

²Department of Urology and ³Physiology, School of Medicine, National Yang-Ming University

Purpose:

Hyperprolactinemia (hyperPRL) is associated with disorder in testosterone (T) release in male rats. The effects contribute to erectile dysfunction and impairment in spermatogenesis. We have confirmed (實際打字時整段須完整，請勿以……表示)

Materials and Methods:

This study we examined the density of autophagy signals in testes of hyperPRL male rats. HyperPRL was induced by alografting of 3 anterior pituitary glands (AP) to subrenal capsule.....

Results:

The data showed that T concentration was much higher in control groups (+CX 280.05 ng/ml) than in hyperPRL groups (+AP 125.32 ng/ml).....

Conclusion:

We confirmed the occurrence of autophagy is much more popular in LC of rats